

Current Biology

Circadian Control Sheds Light on Fungal Bioluminescence

Highlights

- Bioluminescence in *N. gardneri*, a basidiomycete, is regulated by the circadian clock
- Luciferin, reductase, and luciferase, which together make light, all peak at night
- Prosthetic LED-illuminated acrylic mushrooms can be used to study insect behavior
- Insects that can disperse fungal spores are attracted to light at night

Authors

Anderson G. Oliveira,
Cassius V. Stevani, ...,
Jennifer J. Loros, Jay C. Dunlap

Correspondence

stevani@iq.usp.br (C.V.S.),
jay.c.dunlap@dartmouth.edu (J.C.D.)

In Brief

Oliveira et al. report that bioluminescence in *Neonothopanus gardneri* is regulated by the circadian clock. Nighttime light attracts insects that can disperse spores, a benefit to fungi living under the forest canopy, where winds are reduced.



Circadian Control Sheds Light on Fungal Bioluminescence

Anderson G. Oliveira,¹ Cassius V. Stevani,^{2,*} Hans E. Waldenmaier,^{2,3} Vadim Viviani,⁴ Jillian M. Emerson,⁵ Jennifer J. Loros,⁵ and Jay C. Dunlap^{6,*}

¹Departamento de Oceanografia Física, Química, e Geológica, Instituto Oceanográfico, Universidade de São Paulo, São Paulo, SP 05508-120, Brazil

²Departamento de Química Fundamental, Instituto de Química, Universidade de São Paulo, São Paulo, SP 05508-000, Brazil

³Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, São Paulo, SP 05508-000, Brazil

⁴Departamento de Bioquímica, Universidade Federal de São Carlos, Campus Sorocoba, Rodovia João Leme dos Santos, km 110, Sorocaba, SP 18052-780, Brazil

⁵Department of Biochemistry, Geisel School of Medicine at Dartmouth, Hanover, NH 03755, USA

⁶Department of Genetics, Geisel School of Medicine at Dartmouth, Hanover, NH 03755, USA

*Correspondence: stevani@iq.usp.br (C.V.S.), jay.c.dunlap@dartmouth.edu (J.C.D.)

<http://dx.doi.org/10.1016/j.cub.2015.02.021>

SUMMARY

Bioluminescence, the creation and emission of light by organisms, affords insight into the lives of organisms doing it. Luminous living things are widespread and access diverse mechanisms to generate and control luminescence [1–5]. Among the least studied bioluminescent organisms are phylogenetically rare fungi—only 71 species, all within the ~9,000 fungi of the temperate and tropical Agaricales order—are reported from among ~100,000 described fungal species [6, 7]. All require oxygen [8] and energy (NADH or NADPH) for bioluminescence and are reported to emit green light (λ_{max} 530 nm) continuously, implying a metabolic function for bioluminescence, perhaps as a byproduct of oxidative metabolism in lignin degradation. Here, however, we report that bioluminescence from the mycelium of *Neonothopanus gardneri* is controlled by a temperature-compensated circadian clock, the result of cycles in content/activity of the luciferase, reductase, and luciferin that comprise the luminescent system. Because regulation implies an adaptive function for bioluminescence, a controversial question for more than two millennia [8–15], we examined interactions between luminescent fungi and insects [16]. Prosthetic acrylic resin “mushrooms,” internally illuminated by a green LED emitting light similar to the bioluminescence, attract staphilinid rove beetles (coleopterans), as well as hemipterans (true bugs), dipterans (flies), and hymenopterans (wasps and ants), at numbers far greater than dark control traps. Thus, circadian control may optimize energy use for when bioluminescence is most visible, attracting insects that can in turn help in spore dispersal, thereby benefitting fungi growing under the forest canopy, where wind flow is greatly reduced.

RESULTS AND DISCUSSION

Circadian clocks are biological oscillators responsible for maintaining the internal rhythms of animals, plants, fungi, and cyanobacteria to the alternation of external stimuli as light and temperature [17]. Interestingly, bioluminescence and circadian biology share historical antecedents. Studies of the luminescence of the dinoflagellate *Lingulodinium polyedrum* (formerly *Gonyaulax polyedra*), shown to be regulated by a circadian clock, laid the foundations for many of the precepts and paradigms of chronobiology. These include the concept of temperature compensation and the phase response curve, the protocol used to assess the response of the clock to resetting cues as a function of time of day [4]. Although analysis of fungal bioluminescence appeared in the modern literature contemporaneous with that of dinoflagellates [18], the biochemical basis of this luminescence and its possible functional/ecological significance have not been elucidated until very recently [6, 19]. The possibility of circadian regulation was noted in the early 1960s [12, 13], but it was quickly discounted, and in the subsequent 50 years only data questioning rhythmicity have appeared [14, 15].

The Brazilian fungus *Neonothopanus gardneri* belongs to the *Omphalotus* lineage [20] and displays exceptionally intense luminescence from either mycelium or basidiomes (Figures 1A–1D). This fungus is distributed throughout of the Brazilian Coconut Forest (Mata dos Cocais) in the states of Maranhão, Tocantins, Goiás, and Piauí [20], which is a transitional biome between Amazonian Forest and Caatinga. Fruiting bodies grow at the base of palm trees (*Attalea humilis*, *A. funifera*, and *Orbignya phalerata*).

Circadian Control of Bioluminescence

Agar plates freshly inoculated with *N. gardneri* mycelium were maintained over 48 hr in a 12-hr dark/light cycle at constant temperature (21°C, 25°C, and 29°C, depending on the experiment). After this initial entrainment period, the plates were transferred to constant darkness in a climatic chamber equipped with a charge-coupled device (CCD) camera, and an image was digitally recorded every hour over 6 days. Analyses of these pictures




Figure 1. Circadian Control of Fungal Bioluminescence in *Neonothopanus gardneri*

(A–D) Mushroom of *Neonothopanus gardneri* (A) under illumination, and (B) in the dark. The inset shows a rove beetle magnified image. (C) Minimum and (D) maximum bioluminescence from the mycelium of *N. gardneri* during photographic studies of light emission oscillation are shown.

(E) Bioluminescence profile from the mycelium after 48-hr entrainment period at 21°C, 25°C, and 29°C over a 6-day experiment confirms temperature compensation of circadian period length (error bars indicate \pm SD; $n = 3$).

(F) Luminescence profile from luciferin-, reductase-, and luciferase-rich extracts (error bars indicate \pm SD; $n = 3$) obtained from mycelium after 48-hr entrainment period at 25°C and over 48 hr (see more details in the [Experimental Procedures](#)). Luciferin light emission is multiplied by 1,000.

Photos were obtained with a Nikon D3100, equipped with AF-S Micro Nikkor 60 mm f/2.8G ED lenses (mushrooms: ISO 6400, f/5.6, 1/40 s [under headlamp illumination] and 15 s [in the dark]; cultures: ISO 12800, f/4.5, 30 s). Mycelium of *N. gardneri* was cultivated in 2.0% agar medium containing 1.0% sugar cane molasses and 0.10% yeast extract.

by imaging software confirmed that the light emission from the bioluminescent mycelium oscillates in a circadian rhythm of \sim 22 hr at 25°C (Figure 1E), with the peak phase of intensity occurring about 10 hr after the light-to-dark transfer and at circadian intervals thereafter. Circadian oscillators are temperature compensated; i.e., they maintain an \sim 24-hr period whenever the organisms are grown at warmer or cooler temperatures across the physiological range [17]. When the experiments were repeated at 21°C and 29°C, the period was only slightly changed; comparison of the periods at different temperatures allowed calculation (http://www.physiologyweb.com/calculators/q10_calculator.htm) of a Q_{10} of 1.04 for the rhythm, confirming its temperature compensation. Interestingly, these circadian period lengths of less than 24 hr are similar to those reported for *Neurospora crassa*, the non-luminescent ascomycete that is a prominent model system for analysis of the molecular basis of rhythms [21].

Biochemical Basis of Circadian Bioluminescence

Luciferases (enzymes that catalyze the light emission in luminous organisms) have successfully been employed as reporters to monitor a wide array of biological processes, including rhythms.

Fungal bioluminescence depends on four components, namely, a substrate (luciferin), a soluble NAD(P)H-dependent reductase, a membrane-bound oxygenase (luciferase), and oxygen [19]. Active luciferin can be extracted from cultivated mycelium using boiling citrate buffer under argon atmosphere to prevent oxidation. The enzymes can also be partially purified from the cultivated mycelium using cold phosphate buffer, and the reductase and the luciferase subsequently separated by ultracentrifugation because the luciferase is an insoluble membrane protein. Light emission can be obtained in vitro by mixing of the hot extract (luciferin source) and the cold extract (reductase/luciferase source) and finally by addition of NADPH to trigger the reaction [19] (see more details in the [Experimental Procedures](#)). We took advantage of the natural occurrence of bioluminescence in this fungus and used its own luciferase, reductase, and luciferin to investigate the molecular basis of the oscillation in bioluminescence.

Cultures were entrained as above through 2 days of 12-hr dark/light cycles prior to release into constant darkness, and the specific activity of the reductase and luciferase and the concentration of luciferin were measured every 6 hr over




Figure 2. Field Experiments with Acrylic Resin Mushrooms

(A) Acrylic resin mushroom equipped with a green LED and covered with glue under illumination and in the dark.

(B) Traps were attached to the base of juvenile palms of *Attalea speciosa* in *N. gardneri* abundant areas at night.

(C) Emission spectra of LED lights and *N. gardneri*.

(D) Main specimens captured in glue from light traps or controls (LED turned off) over five nights (19 LED traps and 19 controls). See the [Experimental Procedures](#) for additional details. The dipteran flies ($p = 0.00$) and coleopteran beetles ($p = 0.02$) were statistically significant (binomial test, $p < 0.05$), whereas the Hemiptera ($p = 0.06$) and Hymenoptera ($p = 0.09$) were borderline when compared to the control.

48 hr (Figure 1F). The levels of all three biochemical components clearly cycle with a period of about 1 day, reaching maximum intensity about 6 hr after the light-to-dark transfer and at ~22-hr intervals thereafter; the robust 3- to 4-fold amplitude provides a molecular basis for the observed rhythm in luminescence.

Assessing the Adaptive Significance of Fungal Bioluminescence

Altogether, these data show that *N. gardneri* displays a clearly sustained rhythm in bioluminescence with a steady period of about 1 day whose length is temperature compensated and whose phase is set by the entraining cues of a prior light/dark cycle. While these characteristics are sufficient to define the rhythm as circadian, they do not speak to the adaptive significance of rhythmic bioluminescence to the fungus. Bioluminescence per se may fulfill a variety of bio- and ecological functions depending on the luminous organism, including prey attraction, aposematism, illumination, defense, attack, communication, sexual courtship, or as a simple metabolic byproduct [1]. We considered luminescence as a metabolic byproduct to be unlikely given the existence of a specific luciferin and luciferase and instead considered possible significance in light of the biology of *N. gardneri* and fungi in general.

All fungi require help in getting from place to place to colonize new substrates; some achieve this through the use of winds that can carry lightweight spores, and others rely on animals, especially within highly stratified canopy forests, where wind flow can be restricted near to the ground [22]. In the case of arthropods, spore dispersal can occur by the transportation of spores adhered to the body (ectozoochory) or inside the gut of the animal (endozoochory) [23, 24]. Moreover, spore dispersal in canopy forests is greatest at night or early in the morning, when environment humidity and spore germination are highest [22]. Hence, it is reasonable to hypothesize that nocturnal transport of spores by arthropods provides an effective means of dispersal and grants some advantage to fungi, especially in dense forests.

Fungal bioluminescence is too dim to be perceived by animals during the day but is readily perceived at night (even from smaller and less bright mushrooms than *N. gardneri*), so rhythmic control luminescence would be an attractive means to eliminate wasting energy through daytime luminescence. Work from nearly a quarter century ago suggested that insects can be attracted to bioluminescence [16], and indeed, *N. gardneri* basidiomes are often seen infested by staphilinid rove beetles (Figure 1, inset). Given this, we tested the hypothesis that bioluminescence of *N. gardneri* would attract insects capable of dispersing spores. Acrylic resin “mushrooms” were fabricated and equipped with a green light-emitting diode (LED) on the base of stem such that the size, emission spectrum, and intensity of luminescence corresponded to values typical for *N. gardneri* basidiomes (Figure 2); non-luminescent “mushrooms” lacking the LED provided negative controls. When these were placed in the forest habitat of *N. gardneri*, we saw that hemipterans (true bugs), dipterans (flies), hymenopterans (wasps and ants), and other coleopterans, in addition to rove beetles, were captured by the light traps at greater numbers than seen with the control traps (LED turned off). All orders of insects here reported are capable of perceiving green light [25]. Hence, our data are consistent with the hypothesis that nocturnal clock-controlled bioluminescence renders *N. gardneri* more noticeable to insects and thus provides a selective advantage in spore dispersal not afforded to non-luminescent species.

EXPERIMENTAL PROCEDURES

Cultures, Mycelium Cultivation, and Circadian Measurements

Cultures of *N. gardneri* (Mycobank MB519818) were isolated from fruiting bodies collected in the Brazilian Coconut Forest located in the municipality of Altos, Piauí [20]. The mycelium was cultivated on Petri dishes (100 mm diameter) with medium containing 1.0% (w/v) sugar cane molasses (82.2°Bx, Pol 56%) and 0.10% (w/v) yeast extract (Oxoid) in 2.0% (w/v) agar (Oxoid) [26]. The mycelium was inoculated in the center of the Petri dish on a piece of sterilized dialysis cellulose membrane (3 × 3 cm, Sigma-Aldrich) overlying the agar medium. The dialysis membrane permits prompt harvesting of the entire sample without contamination of the culture medium [26]. After inoculation, the cultures were maintained routinely in a climatic chamber (Percival) for 4 days. Different temperatures were used depending on the set

of experiments. Circadian rhythms experiments in which bioluminescence from the mycelium was tracked using a CCD camera were performed at 21°C, 25°C, and 29°C as described below, whereas experiments with time-dependent extraction of luciferin, luciferase, and reductase were carried out at 25°C. The Q_{10} of the rhythm was calculated from the formula below, where T_1 and T_2 are the temperatures at which the rhythm was measured and R_1 and R_2 are the corresponding rates. It should be noted that these are rates, not periods, rate (or frequency) being the inverse of period; periods at 21°C and 29°C were used for the calculation.

$$Q_{10} = \left(\frac{R_2}{R_1} \right)^{\left(\frac{10}{T_2 - T_1} \right)}$$

Partial Purification of Luciferin, Luciferase, and Reductase

So that partially purified luciferin could be obtained, three dialysis membranes containing the mycelium were removed from the medium and cut in small pieces using a scalpel. Luciferin was extracted from this material using a Potter-Elvehjem homogenizer in 1.5 ml of hot (80°C) extraction buffer A (100 mM citrate [Merck; pH 4.0], 1 mM 2-mercaptoethanol [Sigma], and 5 mM EDTA [Sigma]). Afterward, the homogenate was centrifuged at 5,000 × *g* for 5 min at 4°C and the pellet was discarded, whereas the supernatant, the source of luciferin, was kept in ice. The entire process was carried out under ice and argon atmosphere whenever possible.

Partial purification of the enzymes was conducted by similar extraction of mushrooms, but using cold buffer B (100 mM phosphate [pH 7.6], 1 mM 2-mercaptoethanol, and 5 mM EDTA). After the centrifugation, the supernatant containing both the luciferase and the reductase was ultracentrifuged (Hitachi RP50T-2, rotor P50AT2-716) at 198,000 × *g* for 60 min at 4°C. The supernatant contains the reductase and the pellet, the luciferase. The pellet was re-dissolved in 1.5 ml of extraction buffer B. Partially purified luciferin, luciferase (1.0–1.5 mg/ml), and reductase (1.0–1.5 mg/ml) were kept on ice for immediate use or stored on liquid nitrogen. Proteins concentrations were determined using the Bradford assay [27].

Light-Emission Assays

The luminescent reactions were carried out following the procedure described in literature [26]. In summary, each assay was composed by 100 μl of luciferase, 100 μl of reductase, 50 μl of luciferin (all components were partially purified as described in the reference), 50 μl of 1 mg/ml solution of BSA (Sigma-Aldrich), and 50 μl of 70 mM NADPH (nicotinamide adenine dinucleotide phosphate sodium; Sigma-Aldrich) to trigger the reaction. The light emission was recorded during 1 min using a tube luminometer (Berthold, Sirius FB15) at 25°C. All measurements were performed in triplicate.

Circadian Bioluminescence Rhythm of the Fungus *N. gardneri*—Photographical Studies

The cultures were grown under the conditions described above in a 12-hr dark/light cycle over 48 hr in an incubator equipped with a VersArray 1300B liquid-nitrogen-cooled CCD camera system from Princeton Instruments. The camera was housed within a Percival incubator under conditions of constant darkness and temperature (21°C, 25°C, or 29°C) and was controlled with the WinView software (Princeton Instruments). Beginning on the third day, the light/dark cycle was terminated and the cultures were maintained at total darkness for an additional 6 days, during which bioluminescence was monitored. Acquisition times were set to 10 min with a 50 min delay between images (one frame per hour). Data from the image series were extracted with a customized macro ([28]; L.F. Larrondo, A. Stevens-Lagos, V.D. Gooch, J.J.L., and J.C.D., unpublished data) developed for ImageJ (<http://rsbweb.nih.gov/ij/>) and customized with an Excel interface. Although bioluminescence data can be collected from an entire culture (e.g., [28]), differences in periodicity between older and newer regions of a colony (e.g., [29]) can lead to desynchronization and the appearance of rhythm dampening within a colony. To avoid this, the customized Excel macro allowed collection of luminescence from regions of interest, in this case comprising concentric rings surrounding the point of inoculation, analogous to the regions of interest within growth tubes described in [28]. Data were analyzed using a suite of analysis programs designed for analysis of behavioral and molecular cycles including biolumi-

nescence [30]; as recommended [30], chronic trends were first removed to yield a stable baseline; statistically valid rhythmicity was then confirmed by periodogram analysis, and period estimates were derived using maximum entropy spectral analysis (MESA) [30] as reported in Figure 1E.

Circadian Rhythm Effects on Luciferin, Luciferase and Reductase Luminescent Specific Activity

The cultures were cultivated (same conditions described above) in a 12-hr light/dark cycle during 4 days in an incubator at 25°C. On the fifth day, the light/dark cycle was interrupted and the rhythm experiments were initiated. During the 48-hr experiment, the cultures were kept in total darkness. Circadian control of luciferin, luciferase, and reductase was determined by assay of each one of the components separately from extracts every 6 hr, using stock solutions of the other two components. For example, analysis of circadian control of luciferase variation was accomplished by using 24 Petri dishes of mycelium: three dishes for every 6 hr over 48 hr. The luciferin concentration and the reductase activity were maintained constant during the luciferase experiment by using the same batch of frozen pre-purified luciferin (50 μl) and reductase (100 μl) samples in 350 μl total volume assay. Analogous methodology was used for monitoring the luciferin concentration and the reductase activity.

Ecological Studies with Acrylic Resin Mushrooms

Nocturnal collection of insects directly from *N. gardneri* mushrooms and LED light-trap experiments were performed in the Brazilian Coconut Forest (Mata dos Cocais) biome in Fazenda Cana Brava, municipality of Altos, Piauí, Brazil (5°5'39.5" S, 42°23'12.82" W) during the rainy season of March 2013. Acrylic models of *N. gardneri* were equipped with one 530 nm LED and covered with scentless glue (Tangle-Trap Stick Coating, Tanglefoot). Experiments in the field were conducted by setting the voltage knob of the LED to the minimum value so that the light emitted by the acrylic mushrooms approximated that emitted by real mushrooms (see below). Approximately four lighted traps and four control traps (with the LED turned off) were placed at sundown and collected before the sunrise over five nights in forest areas where *N. gardneri* is abundant. The insects captured by the glue were removed from the traps and stored in 70% ethanol after each night. Traps were then cleaned and prepared for reuse. Over five nights of experiments, 19 light traps and 19 control traps (*n* = 19) were used, and a total of 42 insects from light traps and 12 from controls were captured. Sorting and identification of insects was performed with the help of the entomologists Professors Silvio S. Nihei and Sergio Vanin from Instituto de Biologia, Universidade de São Paulo. Total arthropod counts were compared between lighted and control traps with the binomial test for significance, and the difference was considered significant if *p* < 0.05. As the numbers indicate, we collected very few insects in the acrylic mushroom traps, and all collected insects were consumed during the identification process.

Efforts were made to keep the LED light at levels comparable to that emitted by mushrooms, and both would easily be perceived as bright by nocturnal insects. Because the insects are night active, it follows that they must have accommodations to allow them to see at night. In fact, night-active insects, including beetles and ants, undergo dark adaptations that increase the sensitivity of the rhabdom and change the structure of the ommatidia, allowing even diurnal insects to visually navigate at night under extremely low light levels [31, 32], conditions under which bioluminescence by comparison may appear quite bright. However, the absolute amount of “average” bioluminescence in a mushroom in the field has been hard to estimate. Anecdotally, it is well known by the local inhabitants and has been observed by authors of this manuscript (C.V.S., H.E.W., and A.G.O.) that the amount of bioluminescence emitted by *N. gardneri* varies greatly depending, of course, on the size and age of the fruiting body but also on the ambient humidity, optimal conditions being nighttime after a hot day with an afternoon rain storm and a light evening/night breeze. These were the conditions in the forest during the rainy season in March 2013 when the experiments described in the article were performed.

Bioluminescence and LED Emission Spectra

Bioluminescence of *N. gardneri* mushroom and LED emission spectra were recorded using a Hitachi F-4500 spectrofluorometer set at 400 V PMT voltage

and emission slit 1.0 nm (LED) 700 V PMT voltage and emission slit 20 nm (mushroom).

ACKNOWLEDGMENTS

We dedicate this contribution to commemorate the late J. Woodland Hastings, whose 65 years of pioneering research on bioluminescence and circadian biology continue to illuminate all subsequent work. We are indebted to Dr. Ismael Dantas, owner of Cana Brava Farm in Altos, Piauí, Brazil, for the collection of *N. gardneri*. We also thank Professors Silvio S. Nihei and Sérgio Vanin from Instituto de Biologia, Universidade de São Paulo for help with the identification of insects. This work was supported by grants GM34985 to J.C.D., GM083336 to J.J.L. and J.C.D., 2010/11578-5 to A.G.O., and FAPESP 13/16885-1 and NAP-PhotoTech (the USP Research Consortium for Photochemical Technology) to C.V.S.

Received: November 21, 2014

Revised: January 30, 2015

Accepted: February 3, 2015

Published: March 19, 2015

REFERENCES

- Hastings, J.W., and Wilson, T. (2013). *Bioluminescence Living Lights, Lights for Living*. (Harvard University Press).
- Haddock, S.H.D., Moline, M.A., and Case, J.F. (2010). Bioluminescence in the sea. *Annu. Rev. Mar. Sci.* 2, 443–493.
- Widder, E.A. (2010). Bioluminescence in the ocean: origins of biological, chemical, and ecological diversity. *Science* 328, 704–708.
- Hastings, J.W. (2007). The *Gonyaulax* clock at 50: translational control of circadian expression. *Cold Spring Harb. Symp. Quant. Biol.* 72, 141–144.
- Visick, K.L., and Ruby, E.G. (2006). *Vibrio fischeri* and its host: it takes two to tango. *Curr. Opin. Microbiol.* 9, 632–638.
- Stevani, C.V., Oliveira, A.G., Mendes, L.F., Ventura, F.F., Waldenmaier, H.E., Carvalho, R.P., and Pereira, T.A. (2013). Current status of research on fungal bioluminescence: biochemistry and prospects for ecotoxicological application. *Photochem. Photobiol.* 89, 1318–1326.
- Stajich, J.E., Berbee, M.L., Blackwell, M., Hibbett, D.S., James, T.Y., Spatafora, J.W., and Taylor, J.W. (2009). The fungi. *Curr. Biol.* 19, R840–R845.
- Boyle, R. (1666). Observations and tryals about the resemblances and differences between a burning coal and shining wood. *Phil. Trans.* 2, 605–612.
- Aristotle (350 BCE). *De Anima (On the Soul)*, Book II, Part 7, J.A. Smith, translator. <http://classics.mit.edu/Aristotle/soul.2.ii.html>.
- Desjardin, D.E., Oliveira, A.G., and Stevani, C.V. (2008). Fungi bioluminescence revisited. *Photochem. Photobiol. Sci.* 7, 170–182.
- Mendes, L.F., Bastos, E.L., Desjardin, D.E., and Stevani, C.V. (2008). Influence of culture conditions on mycelial growth and bioluminescence of *Gerronema viridilucens*. *FEMS Microbiol. Lett.* 282, 132–139.
- Berliner, M.D. (1961). Diurnal periodicity of luminescence in three basidiomycetes. *Science* 134, 740.
- Calleja, G.B., and Reynolds, G.T. (1970). The oscillatory nature of fungal luminescence. *Trans. Br. Mycol. Soc.* 55, 149–154.
- Mihail, J.D., and Bruhn, J.N. (2007). Dynamics of bioluminescence by *Armillaria gallica*, *A. mellea* and *A. tabescens*. *Mycologia* 99, 341–350.
- Mihail, J.D. (2013). Comparative bioluminescence dynamics among multiple *Armillaria gallica*, *A. mellea*, and *A. tabescens* genets. *Fungal Biol.* 117, 202–210.
- Sivinski, J. (1981). Arthropods attracted to luminous fungi. *Psyche (Stuttg.)* 88, 383–390.
- Dunlap, J.C., Loros, J.J., and DeCoursey, P.J. (2004). *Chronobiology: Biological Timekeeping*. (Sunderland: Sinauer Associates).
- Hastings, J.W., and Sweeney, B.M. (1957). The luminescent reaction in extracts of the marine dinoflagellate, *Gonyaulax polyedra*. *J. Cell. Physiol.* 49, 209–225.
- Oliveira, A.G., and Stevani, C.V. (2009). The enzymatic nature of fungal bioluminescence. *Photochem. Photobiol. Sci.* 8, 1416–1421.
- Capelari, M., Desjardin, D.E., Perry, B.A., Asai, T., and Stevani, C.V. (2011). *Neonothopanus gardneri*: a new combination for a bioluminescent agaric from Brazil. *Mycologia* 103, 1433–1440.
- Baker, C.L., Loros, J.J., and Dunlap, J.C. (2012). The circadian clock of *Neurospora crassa*. *FEMS Microbiol. Rev.* 36, 95–110.
- Shaw, D.C. (2004). Vertical organization of canopy biota. In *Forest Canopies*, Second Edition, M.D. Lowman and H.B. Rinker, eds. (Elsevier Academic Press), pp. 73–101.
- Lilleskov, E.A., and Bruns, T.D. (2005). Spore dispersal of a resupinate ectomycorrhizal fungus, *Tomentella subililacina*, via soil food webs. *Mycologia* 97, 762–769.
- Malloch, D., and Blackwell, M. (1992). Dispersal of fungal diaspores. In *The Fungal Community: Its Organization and Role in the Ecosystem*, Second Edition, G.C. Carroll and D.T. Wicklow, eds. (CRC Press), pp. 147–171.
- Briscoe, A.D., and Chittka, L. (2001). The evolution of color vision in insects. *Annu. Rev. Entomol.* 46, 471–510.
- Oliveira, A.G., Desjardin, D.E., Perry, B.A., and Stevani, C.V. (2012). Evidence that a single bioluminescent system is shared by all known bioluminescent fungal lineages. *Photochem. Photobiol. Sci.* 11, 848–852.
- Bradford, M.M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248–254.
- Larrondo, L.F., Loros, J.J., and Dunlap, J.C. (2012). High-resolution spatiotemporal analysis of gene expression in real time: in vivo analysis of circadian rhythms in *Neurospora crassa* using a FREQUENCY-luciferase translational reporter. *Fungal Genet. Biol.* 49, 681–683.
- Dharmananda, S., and Feldman, J.F. (1979). Spatial distribution of circadian clock phase in aging cultures of *Neurospora crassa*. *Plant Physiol.* 63, 1049–1054.
- Levine, J.D., Funes, P., Dowse, H.B., and Hall, J.C. (2002). Signal analysis of behavioral and molecular cycles. *BMC Neurosci.* 3, 1.
- Meyer-Rochow, V.B. (1972). The eyes of *Creophilus erythrocephalus* F. and *Sartallus signatus* Sharp (Staphylinidae: Coleoptera). Light-, interference-, scanning electron- and transmission electron microscope examinations. *Z. Zellforsch. Mikrosk. Anat.* 133, 59–86.
- Warrant, E., and Dacke, M. (2011). Vision and visual navigation in nocturnal insects. *Annu. Rev. Entomol.* 56, 239–254.